Gestational Diabetes Mellitus Screening: Current Knowledge, Attitudes and Practice



Shastri Motilal

MBBS Hons., DM Lecturer in Family Medicine U.W.I.

Sunday 26th January, 2014

KAP Survey Design

- Sample comprised of 225 physicians
 - 170 PCP's
 - 55 OBGYN

- Survey instrument
 - De novo questionnaire
 - Online survey
 - Email



Results

• 28% response rate (64)

Average age was 38.5 yrs, 60.9% female

• 57.8% of them were PCP's, 34.4% specialists

 78.2% of the sample saw 5-20 antenatal cases weekly

Do you ROUTINELY screen for GDM?

YES!* - 82.8%

- Method used?
 - Urine Testing 1.9%
 - Blood Testing 26.4%
 - Both 71.7%
- When?
 - Booking 69.8%
 - 1st Trimester 32.1%
 - 2nd Trimester 79.2%
 - 3rd Trimester 43.4%

^{*}Neither age, gender, specialty, duration of practice or number of cases seen were predictive of the decision to screen

Which method of blood testing would you use?


O'Sullivan's 50g - 44.2%

• Fasting blood glucose - 48.1%


Glucose Tolerance Testing - 59.6%

• HbA1c - 7.7%


Diagnostic Criteria used by physicians who would do OGTT.



How often should screening be done?



In your practice, how frequent a problem is gestational diabetes?



If you recognized GDM, how would you manage the patient?

• Refer - 76.2%

Manage on my own - 23.8%

 More specialists (52.4%) vs. PCP's (8.1%) would manage GD on their own (P< 0.001).

If you refer, who would you normally refer to?

Referral to:	PCP's (34)	Specialists (10)	P Value
Obstetrician	88.2%	10%	< 0.0001
Physician	2.9%	80%	< 0.0001
Specialty Clinic	20.6%	20%	1.00
Dietitian	23.5%	70%	0.0187


Where would your GDM patient normally be delivered?

• Private sector - 30.2%

Public hospital - 95.2%

 More specialists (61.9%) were more likely to have their patient deliver in the private sector when compared to PCP's. (16.2%)

"All pregnancies should be screened routinely with a standardized national protocol"



How does this data help us?

Low response rate

Small sample



Insightful

Info gained can help build consensus and policy

Summary

 83% of our physicians involved in antenatal care routinely screen for GDM

 94% agree with the need for routine GDM screening with a standardized national protocol!

Acknowledgements

- Questionnaire Pilot testing
 - Dr Darleen Franco and Professor Bassaw

- Questionnaire Development
 - Kevin Yeates (Med Student)

